

Skin Grafted Wound Instructions (Donor Site Glued)

Skin grafting is a surgical procedure where a piece of normal skin is removed from one area (donor site) and transplanted onto a surgical defect (recipient site) to provide restoration of normal skin contour.

MATERIALS NEEDED

Mild (liquid) anti-bacterial soap (Softsoap, Dial, etc)

Tape and Non-stick dressing pads (if covering donor site)

CARE OF SKIN GRAFTED AREA (RECIPIENT SITE)

A special pressure dressing was placed over your skin graft. This was designed to stay in place for 1 week so that you have no wound care for this site for the first week. *NEVER attempt to remove the original bandage.*

1. If this bandage becomes loose on any edges, do not remove any tape. Apply more tape to the outside of the bandage to keep it airtight and maintain pressure. The first week is the most important period for a skin graft to have pressure. If you have any questions call our office.
2. YOU SHOULD NOT GET THIS BANDAGE WET UNTIL YOU RETURN TO OUR OFFICE. For some patients this means no showers. A bath is an alternative for most patients. You can also have someone wash your hair if your site is on the head or neck.
3. You should expect some mild drainage from the wound to stain the edge of the bandage. If you experience excessive drainage, call our office.

CARE OF SUTURED AREA (DONOR SITE)

1. Keep the area dry on the day of surgery. Starting the following day, you may gently clean with a soap and water solution.
2. There should not be excessive drainage from this site. Notify our office if all or the majority of the adhesive comes off at once.
3. Do not swim in pools/hot tubs while the skin adhesive remains on the wound, since it is dissolved by chlorine.
4. For cosmetic purposes or to protect from prolonged sunlight, you may cover your site with a band aid or non-stick dressing and tape, but do not apply anything to the site such as topical antibiotics, lotions, creams, or make up. Oils will cause the skin adhesive to come off prematurely. Also, be sure the tape is not applied too closely to the glued area, as this may cause the skin adhesive to be pulled off when the bandage is changed.
5. If the adhesive begins to peel do not pull it off or pick at it. Please allow it to fall off naturally. This can take up to 2 weeks.
6. Avoid strenuous activities for at least 2 weeks after your surgery.
7. Once the adhesive has completely come off and the wound has healed over, begin using sunscreen to prevent darkening of the scar.

What to Expect:

1. After your surgery you can expect some discomfort from the surgical site. This is highly variable among patients and is also dependent upon the location of your surgery. Some areas are more sensitive than others. We recommend taking a dose of *both* Tylenol 1000mg and ibuprofen 400mg (at the same time) shortly after surgery, and every 6 hours as needed for pain. If your pain is not adequately relieved, or if you are having severe pain then you should call our office immediately or Dr. Viehman after hours. Avoid any aspirin containing products. CAUTION: Bufferin, Anacin, Goody Powders, Excedrin and B.C. Powders all contain aspirin products.
2. After surgery you should expect some mild swelling and redness around the surgical site. In many cases the swelling is a side effect of the numbing medication used during surgery. Some areas of the body tend to swell more than others (around the eyes and on the forehead.) Swelling may worsen over the 48 hours following surgery, particularly following lying flat at night. If you feel your swelling or redness is abnormal please call our office.
3. A low grade fever (99-101) for which any NON-ASPIRIN product may be used. Tylenol is recommended.
4. Mild drainage from the wound may have a foul odor and be yellowish in color. This is especially common on the legs. The yellow drainage is NOT a sign of infection.
5. Significant bleeding is unlikely but can occur. If you experience bleeding then lie down and apply firm, constant pressure to the surgical site for a minimum of twenty minutes. Do not "peek" during this period. If bleeding continues, repeat the pressure on the surgical wound for an additional twenty minutes. If the bleeding persists then please contact our office as early as possible during the day so that we may make arrangements for your evaluation. Dr. Viehman or his PA may also be reached after hours on their cell numbers. If you are unable to reach our office or Dr. Viehman, then please proceed to the nearest emergency room for evaluation and assistance. Call 911 for severe problems.
6. Ice packs may be placed over the wound dressing to help with pain and swelling, and bleeding. The ice pack is placed over the wound for fifteen minutes and may be repeated four times per day. You may also use a bag of frozen peas in substitution.
7. Avoiding/decreasing tobacco use following surgery will drastically improve healing.
8. The donor site will take approximately two weeks to heal. Redness is common and may persist for months. This will eventually resolve. When you return to our office the skin grafted area may vary in color from white to dark purple.

Please contact our office or your local M.D. should you experience excessive bleeding, redness, swelling, fever or pain.

Further questions can be addressed through our office at 910- 256-2100. Our normal business hours are Monday, Tuesday and Thursday from 7:30am – 5:00pm and 7:30am – 1:00pm on Wednesday and Friday.